

# UV Radiation as an Attractor for Insects

Alessandro Barghini PhD<sup>1\*</sup> and Bruno Augusto Souza de Medeiros<sup>2</sup>

*Abstract—Light pollution due to exterior lighting is a rising concern. While glare, light trespass and general light pollution have been well described, there are few reported studies on the impact of light pollution on insects. By studying insect behavior in relation to artificial lighting, we suggest that control of the UV component of artificial lighting can significantly reduce its attractiveness, offering a strong ability to control the impact on insects. Traditionally, the attractiveness of a lamp to insects is calculated using the luminous efficiency spectrum of insect rhodopsin. This has enabled the development of lamps that emit radiation with wavelengths that are less visible to insects (that is, yellow lamps). We tested the assumption that the degree of visibility of a lamp to insects can predict its attractiveness by means of experimental collections. We found that the expected lamp's visibility is indeed related to the extent to which it attracts insects. However, the number of insects attracted to a lamp is disproportionately affected by the emission of ultraviolet radiation. UV triggers the behavior of approaching lights more or less independently of the amount of UV radiation emitted. Thus, even small amounts of UV should be controlled in order to develop bug-free lamps.*

*Keywords—UV radiation, insects, light pollution, light trespass, environment.*

## 1 INTRODUCTION

Today, light pollution is a relevant issue in the subject of exterior lighting. The issue was first raised by astronomers; light pollution impairs astronomical observations and deprives people of the pleasure of contemplating the dark sky (the International Dark-Sky Association can be consulted for information on astronomical light pollution: [www.darksky.org](http://www.darksky.org)). A more recent concern is the impact of light pollution on ecosystems [Longcore and Rich 2004]. It has been

<sup>1</sup>Laboratório de Estudos Evolutivos Humanos, Instituto de Biociência, Universidade de São Paulo and Instituto de Eletrotécnica e Energia, Universidade de São Paulo; <sup>2</sup>Departamento de Zoologia, Universidade de São Paulo and Department of Organismic & Evolutionary Biology, Harvard University

\*Corresponding author: Alessandro Barghini, E-mail: [barghini@iee.usp.br](mailto:barghini@iee.usp.br)

©2012 The Illuminating Engineering Society of North America  
doi: 10.1582/LEUKOS.2012.09.01.003

shown that light pollution alters oviposition behavior in turtles, the trajectory of migratory birds, and the behavior of small mammals; additionally, light pollution has a strong impact on insects, as is stressed in the ample collective work edited by Rich and Longcore [2006]. These concerns have been addressed in a number of official reports and recommendations [Health Council of the Netherlands 2000; The Royal Commission on Environmental Pollution 2009; Huseynov 2010; IDA and IES 2011] and by some power companies with respect to the design of exterior lighting projects. For example, the Florida Power Company developed a manual for ecological lighting of the seacoast to protect sea turtles [Ernest and Martin 1998].

Although concerns about the effects of lighting on insect populations are quite recent, it has long been known that light attracts insects. Entomologists have spent years developing and perfecting light traps for epidemiological and agricultural surveys [see, for example: Hinton 1974; Szentkirályi 2002], but only in the last decade have studies begun to focus on the attraction potential of regular streetlights and its consequences [see, for example: Scheibe 1999; Kolligs 2000; Eisenbeis and Hänel 2009]. There are reasons to believe that night lighting has a significant effect on insect populations. Eisenbeis [2006], for example, estimated that the streetlights of a 240,000-inhabitant town in Germany may kill approximately 360 million insects per season. Considering that insects serve as pollinators for plants and food for a variety of other animals, this increased mortality could have broader effects. The attraction of insects to lights could also have effects on human health because it could provide a novel means of contact between human populations and disease vectors [Barghini and de Medeiros 2010].

To improve regulations and develop minimum-impact lamps, it is important to understand the causes of insect attraction to lights. Although the precise mechanisms are still controversial [see, for example: D'Arcy Thompson 1917; Verheijen 1958; Baker and Sadovy 1978; Janzen 1983; Nowinszky 2003], it has long been known that a key component that determines the attractiveness of a light source to insects is an emission spectrum ranging between ultraviolet and blue [Dethier 1963; Hollingsworth and others 1968; Mazokhin-Porshnyakov 1969; Mikkola 1972; Hinton 1974; Blomberg and others 1976; Walker and Galbreath 1979; Worth 1979; Rea 1993; Service 1993; van Langevelde and others 2011]. Variability in attraction behavior, however, exists because attraction also depends on a number of other factors, one of which is the insects' main activity phase during the day [Rea 1993]. Diurnal insects are the least affected by light, but they may fly towards illuminated areas or UV lamps when disturbed [Lewontin 1959]. This is likely because such areas are presumed by the insect to be open areas into which it is suitable to fly. For nocturnal insects, attraction to light seems to result from navigational errors [Darcy Thompson 1917; Verheijen 1958; Mazokhin-Porshnyakov 1969; Nowinszky 2003]. During the night, insects navigate using celestial references. By keeping a constant angle to such a reference, the insect can fly in a straight path. If the reference happens to be a terrestrial light source, keeping a constant angle would result in an equiangular spiral path towards the light source. Because UV-green or UV-blue contrasts can be used to distinguish between celestial and terrestrial objects [Möller 2002], ultraviolet radiation is probably essential for a light source to be considered as a celestial reference or open space.

Based on this model, strategies to minimize insect attraction to lights are usually based on the spectral responses of insect photoreceptors. The 8<sup>th</sup> edition of the IES Handbook [Rea 1993], for example, follows the recommendation of Barrett and others, [1973, 1974] in suggesting the "maximum use of yellow-red light and the reduction of ultraviolet and blue" (p. 156), avoiding metal fixtures that may reflect

polarized light, the use of directional fixture, and the suggestion that “an attracting lamp can be shaded so that its radiant output is directed downward and confined to [the] immediate area (p. 157)”. Although the IES Handbook did not address this issue in later editions, this model is still generally followed in the design of “bug-free” lamps. Such lamps are usually designed to emit yellowish light because this is the region of the light spectrum that is least visible to insects. Similarly, electric fly killers can be enhanced with a UV-emitting lamp. In both cases, it is implicitly assumed that insects are attracted to lamps that are more visible to them. A recent report in which this assumption is also made is that of van Langevelde and others, [2011]; these authors correlated the mean lamp wavelength with the abundance and diversity of moths attracted as well as with their eye size.

More accurate quantification of the visibility of a lamp to an insect should take insect spectral sensitivity into account, but yellow bug-free lamps are indeed both less visible to insects and have a higher mean wavelength than UV-radiating insect attraction lamps. However, the attraction of a particular lamp for insects is not necessarily related to its visibility or to the mean wavelength it emits. Insects possess a variety of photoreceptors that are not used exclusively for color vision. Some stereotyped behavioral sequences, called wavelength-selective behaviors, are activated by a particular wavelength of light [Goldsmith 1990, 1994]. As reported by Goldsmith [1994:302], “The butterfly *Pieris* exhibits several different behavioral responses to colored lights, each with a distinct action spectrum exhibiting maximum sensitivity at different wavelengths: escape ( $A_{\max}$  370 nm), feeding ( $A_{\max}$  450 nm with a secondary maximum at 600 nm), drumming ( $A_{\max}$  560 nm), and egg laying ( $A_{\max}$  540 nm) [Scherer and Kolb, 1987]. Most of the spectral sensitivity curves are narrower than the absorption spectra of visual pigments, and with mixtures of 600 and 558 nm light, both feeding and drumming are inhibited by the presence of inappropriate wavelengths. The neural wiring thus appears to be more complicated than if each behavior were driven by a single spectral type of receptor.” When perceived by an insect, UVA radiation could trigger a wavelength-dependent response to light attraction similar to what has been measured in frogs: “If the tendency of most species of frogs to jump towards a light is measured as in a forced choice experiment, short wavelengths ( $A_{\max}$  480 nm) stimulate positive phototaxis and longer wavelengths inhibit [phototaxis].” [Goldsmith 1994:303]. There is evidence that this does in fact occur. Insects become disoriented and less active inside greenhouses covered by UV-blocking polyethylene [Antignus 2000]. Moreover, when mulch (a protective cover placed over the soil to retain moisture, reduce erosion, provide nutrients, and suppress weed growth and seed germination) reflects UV radiation, there is a reduction in the population of insect pests [Kring and Schuster 1992]. In the former situation, the absence of UV radiation may disorient insects by creating a “skyless” environment so that an insect would not know where it is able to fly. In the latter case, UV reflection from below would result in an environment with “too much sky,” that is, the insect’s perception of too much space in which it is able to fly. Both of these examples indicate that UV radiation is used by insects to navigate while flying.

Based on the data presented above, it is generally accepted that UV radiation attracts insects. However, this apparent attraction could result from two distinct mechanisms. First, UV radiation might not have any special meaning to the animals and might be attractive only because most insects have a high sensitivity to light in this wavelength range (that is, UV radiation makes lamps more visible to insects). Alternatively, UV radiation may trigger wavelength-selective behavior that results in attraction to the light. It is important to distinguish

between the two mechanisms. If the former holds, a reduction in light attraction would be achieved by reducing lamp radiation on all wavelengths to which insects are most sensitive. If the latter is more important, however, one should eliminate even the smallest amount of UV radiation in order to reduce attraction, and other wavelengths would be less important.

There have been no experimental studies that clearly distinguish the visibility to insects and the UV emission of a lamp while evaluating its attractiveness to insects. Our study aims to test whether UV radiation has a greater attractive power to insects than would be expected from its visibility alone.

## 2 MATERIALS AND METHODS

The test was conducted in a street surrounded by trees and isolated from urban lighting on the “Cidade Universitária” campus of the University of São Paulo. Static insect collecting traps similar to those used by Eisenbeis & Hassel [2000] were set up below lamps installed on seven-meter-tall lampposts.

Each treatment utilized a full cutoff lighting fixture as follows: Hg: 125 w mercury vapor bulb protected with tempered glass; Na: 70 w high-pressure sodium vapor bulb with tempered glass; Hg\_F: 125 w mercury vapor bulb with tempered glass and a UV filter (Polycarbonate Lexan® 2 mm); Na\_F: 70 w sodium vapor bulb with tempered glass and a UV filter (Polycarbonate Lexan® 2 mm); and T: trap without lamp, as a control setup. The radiance spectra of the bulbs were measured with a Monochromator Optronic 740A, an automatic wavelength drive (Optronic 740-1C) and a spectroradiometer (Photo Research OLISA-670). The transmittance of the UV filter was measured with a Hitachi U-3000 spectrophotometer.


The radiance spectrum of each treatment was calculated by multiplying the lamp irradiance by the filter transmittance. The visibility of each treatment to humans and insects was calculated by integrating the treatment radiance after multiplying it by the luminous efficiency spectra of the human eye (photopic vision) and of the rhodopsins of three-rhodopsin insects (represented by the sensitivity curve of *Apis mellifera*).

The collections were performed in two separate campaigns. The first used the Hg, Na, Na\_F and T treatments and totaled 24 collections between March and June 2005; the second used all treatments and totaled 14 collections between March and April 2006. On each collection date, traps were set up before twilight and taken down the following morning. The collected insects were counted and identified to the order level. Ant and termite alates were discarded because a single nest in the surrounding area could significantly bias the results.

The mean insect counts were compared among treatments for both campaigns. The role of UV radiation in the treatments' attractiveness was further tested by fitting the data to a generalized linear model using the visibility to insects, the date of collection and the presence of a UV filter as predictors of insect counts. Specifically, we tested whether accounting for the UV filter significantly improved the fit of the model or whether the treatment visibility was sufficient to explain its attractiveness. This model was adjusted only in the second collecting campaign, in which all treatments were used.

## 3 RESULTS

The number of insects collected varied greatly between collection dates, probably due to meteorological conditions and the lunar phase. Nevertheless, the number of collected insects was clearly higher in the Hg treatment than in the T



**Fig. 1.**  
Mean values and 95 percent confidence intervals of the number of insects collected in each treatment.

treatment (Fig. 1). The same pattern was found for most insect orders when analyzed separately (Table 1).

Hg lamps have a strong UV component, a shorter mean wavelength and a higher visibility to insects than the other lamps tested. In contrast, only a tiny fraction of the emission of a Na lamp is in the UV range, and Na lamps are more visible to humans than Hg lamps but less visible to insects. The use of a UV filter only slightly affects the average wavelength of Na and Hg lamps or their visibility to humans but has a strong effect on the lamps' visibility to insects. The only lamp that has a UV/Green contrast similar to celestial objects is a Hg lamp; all others fall within the range of terrestrial objects as measured by Möller [2002] (Fig. 2, Table 2).

When the mean number of collected insects was considered with respect to the visibility of each treatment to insects, it became clear that these quantities are not entirely correlated. Specifically, treatments with UV filters collected fewer insects than would be expected from their visibility alone (Fig. 3). Indeed, a model that accounts for UV radiation fits the data significantly better than a model that ignores this variable (Table 3).


#### 4 DISCUSSION AND CONCLUSIONS

Overall, our results confirm the general expectation from models currently in use by the illumination industry and suggest that “yellow” lamps are less attractive to insects than “white” lamps. This pattern was observed for most insect orders, with Coleoptera (beetles) and Diptera (flies and mosquitoes) being the most attracted to all treatments. Although our collections were performed in

Order	Hg	Na	Hg_F	Na_F	T
Diptera	25 ± 5.5	16.2 ± 4.4	9.6 ± 2.9	7.2 ± 2.7	1.9 ± 1.0
Coleoptera	18.4 ± 4.9	8.4 ± 2.7	5.3 ± 1.9	4.2 ± 2.0	1.9 ± 0.7
Hymenoptera	10.3 ± 3.2	6.2 ± 3.8	2.7 ± 1.4	3.1 ± 2.1	0.1 ± 0.1
Hemiptera	6.6 ± 2.1	4.3 ± 1.3	1.4 ± 0.7	2.9 ± 0.6	0.2 ± 0.1
Thysanoptera	5.6 ± 4.2	1.2 ± 0.6	0.6 ± 0.4	1.0 ± 0.4	1.6 ± 0.7
Lepidoptera	4.9 ± 1.7	1.1 ± 0.4	1.3 ± 0.4	0.6 ± 0.3	0.03 ± 0.05
Psocoptera	1.3 ± 0.5	1.2 ± 0.6	0.6 ± 0.3	0.7 ± 0.3	0.1 ± 0.1
Other	0.4 ± 0.2	0.2 ± 0.2	0.2 ± 0.1	0.2 ± 0.1	0.0 ± 0.0

**TABLE 1.**  
Mean ± 95% Confidence Interval of the Insect Counts in Each Treatment, Separated by Order. Hymenoptera does not include ants. “Other” includes the following orders: Blattodea, Dermaptera, Neuroptera, Orthoptera, Trichoptera and Strepsiptera

**Fig. 2.**  
Irradiance spectra of the two kinds of lamp used in the test, the transmittance spectrum of the UV filter and the visual sensitivity curves used in the calculations. The horizontal axis is the wavelength (nm) for all curves.






Treatment	Visibility to Humans	Visibility to Insects	Average Wavelength (nm)	UV/Green Contrast
Hg	1	1	531	49.2
Na	1.33	0.44	607	-21.3
Hg_F	0.88	0.59	564	-37
Na_F	1.17	0.38	609	-19.7

**TABLE 2.** Metrics Calculated from Radiance Spectra. Visibility is calculated relative to Hg. The average wavelength follows the calculation by Van Langevelde et al. (2011). The UV/Green contrast follows Möller (2002, Fig. 3). Negative values are usually found in terrestrial objects, while positive values are found in celestial objects

a tropical environment, a similar faunal composition was obtained in Germany [Ensebeis and Hassell 2000], thereby confirming the general applicability of the results. Additionally, we collected a very low number of moths (Lepidoptera), which are probably the most studied organisms in terms of attraction to light [see, for example: Mikkola 1972; Baker and Sadovy 1978; Worth and Muller 1979]. This could be a result of a bias in our traps toward smaller insects or of an already depleted moth fauna in the highly illuminated city of São Paulo. However, the taxonomic composition of our collections suggests that beetles and flies deserve more attention in studies of attraction to lights. Detailed data on insects of the different taxa that were collected available in the appendix of the doctoral thesis of the senior author of this article [Barghini 2008].

Numerous previous studies have found that light sources with higher wavelengths attract fewer insects, but none of these studies have compared the visibility of different lights to insects. Even if insect spectral sensitivity is taken into account, visibility alone is insufficient to explain the attractiveness of a lamp. As an example, the Hg F lamp is a white lamp that is more visible to insects than a Na lamp, yet it exhibited less attractiveness.

For both lamps tested, the use of a UV filter significantly reduced the number of insects despite the variation in UV content of the light emitted by the lamps. UV radiance is approximately 2 percent of the visible light radiance in sodium vapor lamps and 10 percent in mercury vapor lamps. Therefore, the striking effect found in both cases when a UV filter was used indicates that insect attraction does not depend only on the UV amount and lamp visibility. Even small amounts of UV radiation seem to be sufficient for an object to be identified as celestial, resulting in attraction. UV radiation, therefore, acts as a releaser for



**Fig. 3.** Number of insects collected and the calculated visibility to insects (relative to Hg) for each treatment.

**TABLE 3.**  
**Model Comparison by Analysis**  
**of Deviance. The full model**  
**contains the date, visibility to**  
**insects and the presence or**  
**absence of a UV filter as**  
**predictors of insect counts.**  
**The reduced model includes**  
**only the date and visibility.**  
**The P value was determined by**  
**a chi-square test**

	Resid. Df	Resid. Dev.	Df. Deviance	P value
Full model	54	299.63		
Reduced model	55	398.54	-98.91	<0.001

insects, generating an attraction to artificial lights that is greater than would be expected based on the lamp's visibility alone. Our results fail to support the hypothesis that the presence of UV/Green contrast is fundamental for the recognition of a celestial object by an insect. Using the threshold in UV-green contrast found by Möller [2002], Hg lamps would indeed be considered celestial objects, but Na lamps would be classified as terrestrial. At least for nocturnal insects, the absolute emission of UV above a threshold may be more important than its contrast to other colors in triggering attraction behavior.

Our findings provide an important tool for the design of minimum impact lighting systems. As such, our findings support those of Eisenbeins [2006], who advocated the use of UV filters for streetlights. It is highly advisable that studies on lamp attractiveness to insects take into account not only the lamp's visibility and average wavelength but also the lamps' radiance of even minimal amounts of UV radiation. The wavelength threshold that activates insect attraction behavior remains to be identified. While we have found that reducing UV emission to below 400 nm is effective, it is possible that reducing the emission to below a higher wavelength [for example, 480 nm] may have an even greater effect with minimal consequences for human vision. Studies that take this information into account will enable the lighting industry to develop both environmentally friendly and highly effective lighting systems.

## REFERENCES

- Antignus Y. 2000. Manipulation of wavelength-dependent behaviour of insects: an IPM tool to impede insects and restrict epidemics of insect-borne viruses. *Virus Res.* 71:213-220.
- Baker R, Sadovy Y. 1978. The distance and nature of the light-trap response of moths. *Nature.* 276:818-821.
- Barghini A, de Medeiros BAS. 2010. Artificial Lighting as a Vector Attractant and Cause of Disease Diffusion. *Environ Health Persp.* 118:1503-1506.
- Barghini A. 2008 Influência da iluminação artificial sobre a vida silvestre: técnicas para minimizar os impactos, com especial enfoque sobre os insetos [Influence of artificial lighting over the wildlife: techniques for minimizing impacts, with special attention to insects]. Doctoral Thesis. Instituto de Biociências, Universidade de São Paulo, São Paulo, Brazil. Available at: <http://www.teses.usp.br/teses/disponiveis/41/41134/tde-13062008-100639/>. [Accessed: April 1, 2012].
- Barrett JR Jr., Huber RT, Harwood FW. 1973. Selection of lamps for minimal insect attraction. *Trans Am Soc Ag Eng.* 17:710-711.
- Barrett JR Jr., Killough RA, Hartsok JG. 1974. Reducing insect problems in lighted areas. *Trans Am Soc Ag Eng.* 18:329-30.
- Blomberg O, Itamies J, Kuusela K. 1976. Insect catches in a blended and a black light-trap in northern Finland. *Oikos* 27:57-63.
- D'Arcy Thompson. 1917. On growth and form. Reprint, 1992. Cambridge (UK): Cambridge University Press. 345 p.



Dethier VG. 1963. The physiology of insect senses. New York (NY): John Wiley & Sons Inc. 266 p.

Eisenbeis G, Hänel A. 2009. Light pollution and the impact of artificial night lighting on insects. In McDonnell MJ, Hahs AK, Breuste J, editors. Ecology of cities and towns: A comparative approach. Cambridge (UK): Cambridge University Press. pp. 243–263.

Eisenbeis G, Hassel F. 2000. Zur Anziehung nachaktiver Insekten durch Strassenlanternen - eine Studie kommunaler Beleuchtungseinrichtungen in der Agrarlandschaft Rheinhessens [Attraction of nocturnal insects to street lights – a study of municipal lighting systems in a rural area of Rheinhessen]. *Nat Landsch* 75:145–156.

Eisenbeis G. 2006. Artificial night lighting and insects: Attraction of insects to streetlamps in a rural setting in Germany. In Rich C, Longcore T, editors. 2006. Ecological consequences of artificial night lighting. Washington (DC): Island Press. pp. 281–304.

Ernest RG, Martin RE. 1998. Coastal roadway lighting manual: A handbook of practical guidelines for managing street lighting to minimize impacts to sea turtles. Jensen Beach (FL): Ecological Associates, Inc. 71 p.

Goldsmith TH. 1994. Ultraviolet receptors and color vision: Evolutionary implications and the dissonance paradigm. *Vision Res.* 30:1479–87.

Goldsmith TH. 1990. Optimization, constraint and history in the evolution of eyes. *Q Rev Biol.* 65:281–322.

Health Council of the Netherlands. 2000. Publication no. 2000/25E: Impact of outdoor lighting on man and nature. The Hague (Netherlands): Health Council of the Netherlands. 46 p.

Hienton TE. 1974. Summary of investigations of electric insect traps. Technical Bulletin No. 1498. Washington (DC): Agricultural Research Service. US Department of Agriculture. 136 p.

Hollingsworth JPA, Hartstack Jr. W, Lindquist DA. 1968. Influence of near-ultraviolet output of attractant lamps on catches of insects by light traps. *J Econ Entomol.* 61:515–521.

Huseynov, R. 2010. Noise and light pollution. Doc. 12179. Report of the Committee on the Environment, Agriculture and Local and Regional Affairs. Council of Europe. Available at <http://assembly.coe.int/Main.asp?link=/Documents/WorkingDocs/Doc10/EDOC12179.htm>. [Accessed: April 1, 2012].

[IDA and IES] International Dark-sky Association and Illuminating Engineering Society. 2011. Model Lighting Ordinance, with User's Guide. Available at [http://www.darksky.org/index.php?option=com\\_content&view=article&id=622](http://www.darksky.org/index.php?option=com_content&view=article&id=622). [Accessed: April 1, 2012].

Janzen DH. 1983. Insects. In: Janzen DH, editor. Costa Rican Natural History. Chicago (IL): University of Chicago Press. pp. 619–780.

Kolligs D. 2000. Ökologische Auswirkungen künstlicher Lichtquellen auf nachtaktive Insekten, insbesondere Schmetterlinge (Lepidoptera) [Ecological effects of artificial light sources on nocturnally active insects, in particular on moths (Lepidoptera)]. *Faun-Oekol Mitt Suppl.* 28:1–136.

Kring J, Schuster DJ. 1992. Management of insects on pepper and tomato with UV-reflective mulches. *Fla Entomol.* 75:119–129.

Lewontin R. 1959. On the anomalous response of *Drosophila pseudoobscura* to light. *Am Nat.* 93:321–328.

Longcore T, Rich C. 2004. Ecological light pollution. *Front Ecol Environ.* 2:191–198.

Mazokhin-Purshnyakof G. 1969. Insect vision. New York (NY): Plenum Press. 306 p.

Mikkola K. 1972. Behavioural and electrophysiological responses of night-flying insects, especially Lepidoptera, to near-ultraviolet and visible light. *Ann Zool Fenn.* 9:225–254.

Möller R. 2002. Insects could exploit UV-green contrast for landmark navigation. *J Theor Biol.* 214:619–31.

Nowinszky L. 2003. The orientation of insects by light—major theories. In: The handbook of light trapping. Szombathely (Hungary): Savaria University Press. pp. 15–18.

Rea, MS, editor. 1993. Lighting handbook: Reference & application. 8th ed. New York (NY): Illuminating Engineering Society of North America. 989 p.

Rich C, Longcore T, editors 2006. Ecological consequences of artificial night lighting. Washington (DC): Island Press. 458 p.

Scheibe MA. 1999. Über die Attraktivität von Straßenbeleuchtungen auf Insekten aus nahegelegenen Gewässern unter Berücksichtigung unterschiedlicher UV-Emission der Lampen [On the attractiveness of roadway lighting to insects from nearby waters with consideration of the different UV-emission of the lamps]. *Natur und Landschaft*. 74:144–146.

Service MW. 1993. Mosquito ecology: Field sampling methods. 2nd ed. London (UK): Elsevier. 988 p.

Szentkirályi F 2002. Fifty years-long insect survey in hungary: T. Jeremy's contribution to light-trapping. *Acta Zool Acad Sci Hung*. 48 (supl. 1): 85–105.

The Royal Commission on Environmental Pollution. 2009. Artificial light in the environment. London (UK): The Stationery Office Limited. 44 p.

van Langevelde F, Ettema JA, Donners M, WallisDeVries MF, Groenendijk D. 2011. Effect of spectral composition of artificial light on the attraction of moths. *Biol Conserv*. 144:2274–2281.

Verheijen FJ. 1958. The mechanisms of the trapping effect of artificial light sources upon animals. *Arch Neerl Zool*. 13.

Walker AK, Galbreath RA. 1979. Collecting insects at lights: a test of four types of lamp. *New Zeal Entomol*. 7:83–85.

Worth CB, Muller J. 1979. Captures of large moths by an ultraviolet light trap. *J Lepid Soc* 33:261–264.